

# Changes in the Cell Surface Markers During Normal Hematopoiesis: A Guide to Cell Isolation

Armin Attar<sup>1,2,\*</sup>

<sup>1</sup>Cell and Molecular Research Club, <sup>2</sup>Student Research Committee, Shiraz University of Medical Sciences, Shiraz, Iran

**Abstract:** Hematopoiesis, the process of hematopoietic cell production, largely takes place in the bone marrow (BM) of humans. This process follows a stepwise manner in which hematopoietic stem cells give rise to progenitor cells and they develop the terminally differentiated cells along each lineage through a sequential series of stages. Consequently, constant changes would occur in the gene expressions leading to morphological or functional changes necessary for different stages of maturation. These changes provide us with guides to differentiate different subsets of hematopoietic hierarchy based on the cell surface antigen markers and will help us to isolate various cells from the hematopoietic hierarchy. Here we have a short review on the changes of these surface markers during different stages of development and we have applied an algorithmic approach for the isolation of all these cells based on our current understandings of this system.

**Keywords:** Cell surface antigen markers, Hematopoiesis, Progenitor cell, Stem cell.

## INTRODUCTION

In humans, the process of hematopoietic cell production (Hematopoiesis) occurs largely in the bone marrow (BM). Only the T-cell differentiation occurs outside the BM, principally within the thymus. This process begins with a quiescent stem cell which is stimulated with various environmental factors and gives rise to progenies differentiating along multiple lineages.

The current model for describing the normal hematopoiesis considers a relatively linear fashion for differentiation along each lineage through a sequential series of stages. The final products are fully functional mature cells which are needed for the performance of routine tasks of the hematopoietic system. The maturation of hematopoietic cells along each lineage is tightly controlled by sequential expression of genes and their products [1]. Consequently, the pattern of genes expression and their resultant products shows constant and reproducible changes during morphological or functional stages of maturation.

A group of these proteins are located on the cell surface. They may have diverse functions such as serving as growth factor receptors, adhesion molecules, extracellular enzymes, and signal transduction molecules. Many of these cell surface proteins were simply identified by their antigenic reactivity and the functions for some of these molecules is not clearly

known. Some of these cell-surface antigens are restricted to a specific cell lineage, while others may be more widely expressed but their levels of expression varies with maturational stage or activation state. This may aid us in identification and isolation of different cells within diverse maturational stages based on their patterns of cell surface antigen expression [2].


## HEMATOPOIETIC STEM AND PROGENITOR CELLS

Both repair of the injured tissues and formation of new tissues are dependent on the quiescent stem cells localized within specific niches in each organ [3]. Activation of stem cells is closely controlled by environmental stimulations by the surrounding microenvironment and follows a sequence of events at the genetic and epigenetic levels [4]. This tight controlling machine leads to establishment of a hierarchy starting from the slow-cycling stem cell with subsequent transiently amplifying precursors with a high proliferative capacity, but with lineage-commitment properties. At the end of this spectrum are the non-cycling, terminally differentiated cells that carry on the organ functions [5].

During the development of terminally differentiated cells, the stem cells themselves remain undifferentiated, which passes through their unique capacity for self-renewal. When a stem cell divides into two fates, only one of the resulting daughter cells enters the path for increasing specificity and the other one instead remains undifferentiated [6]. This phenomenon keeps the numbers in the overall stem cell pool constant, whereas the proliferation of the descendant progenitors can rapidly provide populations of specific cell types in

\*Address Correspondence to this author at the Department of Cardiovascular Medicine, School of Medicine, Zand Street, Shiraz University of Medical Sciences, Postal Code: 71344-1864, Shiraz, Iran; Tel: +989177141797; Fax: +987112349521; E-mail: attarar@sums.ac.ir

response to changing needs. The capacity of stem cells to recreate themselves is called self-renewality and is one of the most important defining properties of stem cells. This gives them a potential for a nearly unlimited life span. Progenitor cells do have some ability to renew themselves as well, but their ability to reproduce themselves is restricted by an internal counting mechanism and only finite rounds of duplication can be preceded. The ability of the progenitors' offspring to multiply declines steadily as differentiation goes on (Figure 1). Hematopoietic stem cells (HSCs) are defined as single cells with lifelong ability to self-renew as well as differentiate to produce all blood cell lineages (multipotency).



**Figure 1:** When a stem cell divides into two fates, only one of the resulting daughter cells enters the path for increasing specificity and the other one instead remains undifferentiated. This process is called self-renewality. Consequently the numbers of stem cells remains constant, whereas the descendant progenitors will provide the terminally differentiated cells needed for proper tissue functioning. Progenitor cells do have some abilities to renew themselves as well, which is limited by an internal counting mechanism leading to only finite rounds of duplication.

These hematopoietic cells with repopulating activities have been identified within  $CD34^+$  cells [7]. Civin and colleagues in 1981 reported the presence of a marker, at first named as My10, on about 1% to 4% of BM cells which was also present on KG1A leukemia cells and later shown to be  $CD34$  [8]. In 1988, Bernstein and colleagues showed that transplantation of  $CD34^+$  cells can reconstitute hematopoiesis in lethally irradiated baboons [9], and the Thomas group in 1990 performed the first human  $CD34^+$  enriched autologous graft with successful engraftment in a patient with a brain neoplasm [10]. Primitive hematopoietic cells that tend to lose their self-renewal potential start to express higher levels of the  $CD38$  antigen while becoming more committed to differentiate. Therefore, HSC are found within the  $CD34^+CD38^-$  cell fraction [11-13]. In human, two groups of cells can fulfill the definition for HSCs and therefore both can be named as stem cell. But only one can maintain a long-term multilineage reconstitution (LTMR) in allogenic transplants and therefore has been named as long term HSC (LT-HSC). In irradiation-reconstitution experiments, human  $CD34^+CD90^+Lin^-$  cells were the only cells in human adult and fetal marrow, cord blood, fetal liver, and mobilized peripheral blood (MPB) that achieved LTMR [14]. So, LT-HSCs are  $CD34^+CD38^-CD90^+Lin^-$  cells. These cells can give rise to a progeny, with the phenotype of  $CD34^+CD38^-CD90^-Lin^-$ , which has a short-term self-renewality potential for 10 weeks despite the ability to produce all blood lineages. They are called short term HSCs (ST-HSC) [14]. In contrast, in the murine system, LT-HSCs are  $CD34^+CD38^+$  cells and the more committed progenitors are showing a  $CD34^+CD38^-$  phenotype [15, 16]. Therefore,  $CD38$  is a primitive marker in mice and a differentiation marker in humans.

In addition to the mentioned markers, other antigen combinations have been used to discriminate more primitive from more mature  $CD34^+$  cells. Some of the combinations used are  $CD34^+c-kit^+CD38^-CD33^-Rho-123^-$  [17],  $CD34^+HLA-DR^-Lin^-$  [18], or  $CD34^+CD38^-Rho-123^{low}Lin^-$  [19], and  $CD133^+ALDH^{hi}Lin^-$  [20].

The  $CD133$  marker, also known as prominin-1 or AC133 antigen, is also co-expressed on most of the  $CD34^+$  cells, and can characterize stem cells in humans [21-23]. Investigations have shown the presence of very rare undifferentiated cells with stem cell properties within the  $Lin^-CD34^+CD38^-$  cell fraction which expresses  $CD133$  [24-26]. It seems that  $CD133$  describes the most primitive human hematopoietic cells more precisely than  $CD34$  [7].

Table 1: Expression of Surface Antigens during Different Stages of Granulopoiesis (Neutrophilic Maturation)

	GMP	Myeloblast	Promyelocyte	Myelocyte	Metamyelocyte	Band Cell	Neutrophil
CD10	-	-	-	-	-	-	++
CD11a	++	+/++	+/++	+/++	+/++	+/++	+/++
CD11b	-	-	-	+/++	++	++	++
CD11c	-	-	-	++	++	++	++
CD13	++	+/+++	+/+++	+	+/++	++	+++
CD14	NA <sup>b</sup>	-	-	-	-	-	+
CD15	-	+	+++	+++	+++	+++	+++
CD15s	++	+++	+++	++	+	+	+++
CD16	-	-	-	-	+	++	+++
CD18	++	++	+	+++	++	-	-
CD24	-	-	-	++	++	++	++
CD29	+++	++	++	++	+	+	+
CD31	+++	++	++	++	+	+	+
CD32	++	++	++	++	++	++	+++
CD33	+++	+++	+++	++	+	+	+
CD34	+	++	-	-	-	-	-
CD35	-	-	-	-	-	++	++
CD44	+++	+++	++	+	+	++	+++
CD45	NA <sup>a</sup>	+/++	+/++	++	++	++	++
CD45RA	++	+	+	-	-	-	-
CD45RO	-	-	-	+	+	++	+++
CD49d	+++	++	++	++	+	-	-
CD49e	+++	++	++	++	+	+	+
CD54	++	++	++	+/-	+/-	+/-	+/-
CD55	+++	+++	+	+	+++	+++	+++
CD59	+++	+++	+++	+++	+++	+++	+++
CD62L	++	++	++	++	++	++	++
CD64	++	+	+	++	++	-	-
CD65	+	+	++	++	++	+++	+++
CD66a	-	-	-	++	++	++	++
CD66b	-	-	+++	+++	++	++	++
CD82	+++	++	++	++	++	++	++
CD87	-	-	-	-	-	++	++
CD117	NA <sup>a</sup>	++	+	-	-	-	-
CD162	+++	+++	+++	+++	+++	+++	+++

[Modified from Elghetany MT: Surface Antigen Changes during Normal Neutrophilic Development: A Critical Review. Blood Cells Mol Dis 2002; 28: 260-74]. GMP means granulocyte monocyte progenitor cell; +++ means strong expression; ++ means moderate expression; + means weak expression; +/- means very weak expression; - means no expression; NA means Data Not Available; a means possible expression; b means possible not expressing.

Human hematopoietic progenitors have also been characterized. Manz *et al.* described the characteristics of the "Common Myeloid Progenitor Cell (CMP)", "Granulocyte Monocyte Progenitor Cell (GMP)", and

"Megakaryocyte Erythrocyte Progenitor Cell (MEP)" within the CD34<sup>+</sup>CD38<sup>+</sup> fraction of hematopoietic cells from both the adult bone marrow and cord blood by the differential expression of CD45RA and IL3Ra (CD123)

[27]. They showed that CMP (CD123<sup>+</sup>CD45RA<sup>-</sup>) can give rise to both the GMP (CD123<sup>-</sup>CD45RA<sup>-</sup>) and the MEP (CD123<sup>low</sup>CD45RA<sup>+</sup>). None of these cells contained a lymphoid developmental potential [14]. Instead, in 1995, Galy *et al.* described CD34<sup>+</sup>CD38<sup>+</sup>Lin<sup>-</sup>CD10<sup>+</sup>CD45RA<sup>+</sup> cells within human BM that could give rise to T, B, natural killer (NK), and dendritic cells and are common lymphoid progenitor cells [28].

## GRANULOPOIESIS

The process of Granulocyte maturation (Granulopoiesis) starts from myeloblasts through a linear process into morphological stages historically termed promyelocytes, myelocytes, metamyelocytes, bands, and neutrophils. Only the last fates within this lineage (band/neutrophils) are fully functional and have bactericidal potentials. The antigenic changes during these stages have been widely studied from an immunophenotypic perspective. [2].

Among the CD markers, CD15 needs special attention in granulocyte maturation charts. Early studies suggested the expression of CD15 from GMP through the segmentation phase of neutrophils [29]. In other studies, expression of CD15 was shown at the blast stage but not on CD34<sup>+</sup> cells [30-32]. More recent studies indicate that a small portion of CD34<sup>+</sup> bone marrow precursor cells do express CD15 and this expression increases significantly with sequential maturation from promyelocytes to neutrophils [33, 34]. So it seems that expression of CD15 is the only key to differentiate between myeloblast and GMP [35]. A summary of antigen expression during different developmental stages of neutrophils is shown in Table 1.

## MONOCYTE MATURATION

Monocytes maturation starts from monoblasts as a continuum through one intermediate stage termed the promonocyte [32, 36]. With maturation of monoblasts from GMPs, expression of CD4 begins and increases with more differentiation. In contrast, CD34 expression stops with transition from monoblasts toward promonocytes. The expression of CD64, HLA-DR, and CD33 are retained at relatively high levels on maturing monocytes unlike the myeloid counterparts and may serve as useful monocytic markers [2].

In general, the presence of CD4 on monoblast and its progenies can help differentiate them from myelocytic lineages and also GMP. The presence of CD33 makes them different from T-cell subtypes [2]. A

summary of antigen expression during different developmental stages of Monocytes is shown in Table 2.

**Table 2: Expression of Surface Antigens during Different Stages of Monocytic Maturation**

	GMP	Monoblast	Promonocyte	Monocyte
CD4	-	-/+	+	+
CD11b	-	-	++	+++
CD13	++	++	+ / ++	++ / +++
CD14	NA <sup>b</sup>	-	+ / ++	+++
CD15	-	-	++	+
CD16	-	-	-	- / +
CD33	+++	+++	+++	+++
CD34	+ / ++	+	-	-
CD36	-	-	++	+++
CD45	NA <sup>a</sup>	+	++	+++
CD64	++	-	++	+++
HLA-DR	++	++	+++	++ / +++

[Modified from Wood B: Multicolor immunophenotyping: human immune system hematopoiesis. *Methods Cell Biol.* 2004; 75: 559-76.]

GMP means granulocyte monocyte progenitor cell; +++ means strong expression; ++ means moderate expression; + means weak expression; +/- means very weak expression; - means no expression; NA means Unknown Data Not Available; a means possible expression; b means possible not expressing.

## ERYTHROPOIESIS

Erythropoiesis is the process of formation of red blood cells. The maturational stages of this process have historically been divided into proerythroblasts, basophilic erythroblasts, polychromatophilic erythroblasts, and erythrocytes based on their morphological appearance. The morphological changes have shown reasonable correlations with the observed immunophenotypes [37-40]. The expression of CD34 is high on MEPs but low on proerythroblasts. The expression of CD38 also decreases with maturations. Proerythroblasts show high levels of CD71 and CD235a, both of which being absent on MEPs. It should be noted that the preparations of BM samples for evaluating maturational stages during erythropoiesis differ from those for other lineages. For example, usage of lysing reagents such as NH<sub>4</sub>Cl generally leaves only proerythroblasts for evaluation [2]. A summary of antigen expression during different developmental stages of erythrocytes is shown in Table 3.

## MEGAKARYOPOIESIS

Megakaryopoiesis is the process that leads to the production of the platelets. The development of the

**Table 3: Expression of Surface Antigens during Different Stages of Erythropoiesis (Erythrocyte Maturation)**

	MEP	Proerythroblast	Basophilic Erythroblast	Polychromatophilic Erythroblast	Erythrocyte
CD34	++	-/+	-	-	-
CD36	-	++	+++	++	-
CD38	++	+	-/+	-	-
CD45	++	+	-/+	-	-
CD45RA	-	NA	NA	NA	NA
CD71	-	+++	+++	+++	-
CD117	++	++	-	-	-
CD123	-	NA	NA	NA	NA
CD235a	-	+ / ++	+++	+++	+++
HLA-DR	++	++	+	-	-
Hemoglobin	-	-	-/+	+	+++

[Modified from Wood B: Multicolor immunophenotyping: human immune system hematopoiesis. Methods Cell Biol. 2004; 75: 559-76.]

MEP means megakaryocyte–erythrocyte progenitor cell; +++ means strong expression; ++ means moderate expression; + means weak expression; +/- means very weak expression; - means no expression; NA means Data Not Available.

platelet passes through megakaryoblast, promegakaryocytes, and megakaryocytes. Production of the platelets from mature megakaryocytes needs cytoplasmic fragmentation occurring through a dynamic and regulated process, called proplatelet formation. This process consists of pseudopodial elongations with final breaking into the blood.

The hallmarks of this lineage are CD41 and CD61. CD41 is present on about 3% of marrow CD34<sup>+</sup>. These CD34<sup>+</sup>CD41<sup>+</sup> cells are enriched in megakaryoblasts. The expression of CD41 precedes the detection of CD42 [41]. Thus, CD34<sup>+</sup>CD41<sup>+</sup>CD42<sup>-</sup> cells correspond to true megakaryoblast whereas CD41<sup>+</sup>CD42<sup>+</sup> cells give rise to promegakaryocytes and the expression of CD42 accompanies late differentiation steps. CD61 expression starts with passing through MEP to megaka-

ryoblast and can help differentiate these two cells. As the maturation of cells increases, the CD34 antigen disappears [42].

A summary of antigen expression during different developmental stages of megakaryocytes is shown in Table 4.

**CONCLUSION**

Although, to the best of our knowledge, no study has evaluated all the CD markers together simultaneously, the sum of previous investigations can logically be mixed to reach the final picture of the whole hematopoietic hierarchy (Figure 2).


Our current knowledge from the widely popular model of describing the hematopoietic hierarchy, the “sequential determination model”, provides us with guides to isolate different cells within different stages of maturation (Figure 3) and helps us differentiate various subsets of hematopoietic hierarchy based on cell surface antigen markers. However, there are still some missing parts; for instance, the status of CD38 expression on the Myeloblasts and Monoblasts is not determined.

Unlike the classical sequential determination model, two human cord blood derived cell populations (CD34<sup>+</sup>CD45RA<sup>high</sup>CD10<sup>+</sup>Lin<sup>-</sup> and CD34<sup>+</sup>CD45RA<sup>high</sup>CD7<sup>+</sup>Lin<sup>-</sup>) have been described by Haddad and colleagues to contain robust myelomonocytic differentiation potentials despite being primed to

**Table 4: Expression of Surface Antigens during Different Stages of Megakaryopoiesis (Platelet Maturation)**

	MEP	Megakaryoblast	Promegakaryocyte	Megakaryocyte
CD34	++	-/+	-	-
CD38	++	-/+	+	++
CD41	++	+	+	++
CD42	-	-	-/+	+
CD61	-	+	+	++


MEP means megakaryocyte–erythrocyte progenitor cell; +++ means strong expression; ++ means moderate expression; + means weak expression; +/- means very weak expression; - means no expression.



**Figure 2:** A schematic representation for how different cells within hematopoietic hierarchy express various cell surface markers.

develop the T-cell and NK cell or the B-cell lineage, respectively [43]. However, this CD34<sup>+</sup>CD38<sup>-</sup>CD7<sup>+</sup> population does not exist in the adult bone marrow [14]. Similarly, CD34<sup>+</sup>CD10<sup>-</sup>CD19<sup>-</sup> cells are known to be committed to early B-cell development do still have the

capacity to differentiate into NK-cells, T-cells and macrophages [44]. Additionally, CXCR4<sup>-</sup>CD34<sup>+</sup>CD19<sup>+</sup> B-cell progenitor cells have been shown to have potentials for granulocyte, macrophage, and red blood cell development [45].



**Figure 3:** A proposed algorithmic chart for isolation of cells within hematopoietic hierarchy based on the cell surface antigen markers. (Abbreviations: CLP=Common Lymphoid Progenitor cell, CMP = Common Myeloid Progenitor Cell, GMP = Granulocyte Monocyte Progenitor Cell, LT-HSC = Long Term Hematopoietic Stem Cell, MEP = Megakaryocytic Erythroid Progenitor Cell, MN = Mononuclear Cell, ST-HSC = Short Term Hematopoietic Stem Cell.).

The identification of progenitors with diverse cell fates has prompted new models of hematopoiesis [46-50]; however, most of these models lack the consensus for a lineage branching map. A revised model to depict the developmental relationships between hematopoietic cell lineages has been proposed by Ceredig and colleagues. This model involves grouping the pairwise relationships between lineage fates around a broken circle, and considers the

hematopoietic hierarchy as continuum other than a branched stepwise model [51].

Although these new models may be more realistic, because the differentiation of hematopoietic cells is a continuous rather than a step-wise process, they models cannot help us in isolating cells from this spectrum and the classical models seem to be more practical for this purpose.

## ACKNOWLEDGMENT

The author would like to thank Dr. Nasrin Shokrpour at Center for Development of Clinical Research of Nemazee Hospital for editorial assistance.

## REFERENCES

- [1] Payne KJ, Crooks GM. Human hematopoietic lineage commitment. *Immunol Rev* 2002; 187: 48-64.
- [2] Wood B. Multicolor immunophenotyping: human immune system hematopoiesis. *Methods Cell Biol.* 2004; 75: 559-76.
- [3] Rizvi AZ, Wong MH. Epithelial stem cells and their niche: there's no place like home. *Stem Cells* 2005; 23: 150-65.
- [4] Bissell MJ, Labarge MA. Context, tissue plasticity, and cancer: are cancer stem cells also regulated by the microenvironment? *Cancer Cell* 2005; 7: 17-23.
- [5] Attar A, Khosravi Maharloo M, Khoshkhou S, Hosseini A, Jaberipour M, Dehghan A, *et al.* Colony forming unit endothelial cells do not exhibit telomerase alternative splicing variants and activity. *Iran Biomed J* 2013; 17: 146-151.
- [6] National Institute of Health. *Stem Cells: Scientific Progress and Future Research Directions.* University Press of the Pacific Honolulu, Hawaii; 2001.
- [7] Giebel B, Punzel M. Lineage development of hematopoietic stem and progenitor cells. *Biol Chem* 2008; 389: 813-24.
- [8] Civin CI, Strauss LC, Brovall C, Fackler MJ, Schwartz JF, Shaper JH. Antigenic analysis of hematopoiesis: III. a hematopoietic progenitor cell surface antigen defined by a monoclonal antibody raised against KG-1a cells. *J Immunol* 1984; 133: 157-165.
- [9] Berenson RJ, Andrews RG, Bensinger WI, Kalamasz D, Knitter G, Buckner CD, *et al.* Antigen CD34+ marrow cells engraft lethally irradiated baboons. *J Clin Invest.* 1988; 81: 951-955.
- [10] Berenson RJ, Bensinger WI, Hill R, Andrews RG, Garcia-Lopez J, Kalamasz DF, *et al.* Stem cell selection-clinical experience. *Prog Clin Biol Res* 1990; 333: 403-410; discussion 411-413.
- [11] Terstappen LW, Huang S, Safford M, Lansdorp PM, Loken MR. Sequential generations of hematopoietic colonies derived from single nonlineage-committed CD34+CD38- progenitor cells. *Blood* 1991; 77: 1218-1227.
- [12] Hao QL, Thiemann FT, Petersen D, Smogorzewska EM, Crooks GM. Extended long-term culture reveals a highly quiescent and primitive human hematopoietic progenitor population. *Blood* 1996; 88: 3306-3313.
- [13] Laroche A, Vormoor J, Hanenberg H, Wang JC, Bhatia M, Lapidot T, *et al.* Identification of primitive human hematopoietic cells capable of repopulating NOD/SCID mouse bone marrow: implications for gene therapy. *Nat Med* 1996; 2: 1329-1337.
- [14] Weissman IL, Shizuru JA. The origins of the identification and isolation of hematopoietic stem cells, and their capability to induce donor-specific transplantation tolerance and treat autoimmune diseases. *Blood* 2008; 112: 3543-3553
- [15] Osawa M, Hanada K, Hamada H, Nakauchi H. Long-term lymphohematopoietic reconstitution by a single CD34 low/negative hematopoietic stem cell. *Science* 1996; 73: 242-245.
- [16] Randall TD, Lund FE, Howard MC, Weissman IL. Expression of murine CD38 defines a population of long-term reconstituting hematopoietic stem cells. *Blood* 1996; 87: 4057-4067.
- [17] Liu H, Verfaillie CM. Myeloid-lymphoid initiating cells (ML-IC) are highly enriched in the rhodamine-ckit+CD33-CD38- fraction of umbilical cord CD34+ cells. *Exp Hematol* 2002; 30: 582-589.
- [18] Verfaillie C, Blakolmer K, McGlave P. Purified primitive human hematopoietic progenitor cells with longterm *in vitro* repopulating capacity adhere selectively to irradiated bone marrow stroma. *J Exp Med* 1990; 172: 502-509.
- [19] McKenzie JL, Takenaka K, Gan OI, Doedens M, Dick JE. Low rhodamine 123 retention identifies long-term human hematopoietic stem cells within the Lin-CD34+CD38- population. *Blood* 2007; 109: 543-545.
- [20] Hess DA, Wirthlin L, Craft TP, Herrbrich PE, Hohm SA, Lahey R, *et al.* Selection based on CD133 and high aldehyde dehydrogenase activity isolates long-term reconstituting human hematopoietic stem cells. *Blood* 2006; 107: 2162-2169.
- [21] Yin AH, Miraglia S, Zanjani ED, Almeida-Porada G, Ogawa M, Leary AG, *et al.* AC133, a novel marker for human hematopoietic stem and progenitor cells. *Blood* 1997; 90: 5002-5012.
- [22] de Wynter EA, Buck D, Hart C, Heywood R, Coutinho LH, Clayton A, *et al.* CD34+AC133+ cells isolated from cord blood are highly enriched in long-term culture-initiating cells, NOD/SCID-repopulating cells and dendritic cell progenitors. *Stem Cells* 1998; 16: 387-396.
- [23] Giebel B, Corbeil D, Beckmann J, Hohn J, Freund D, Giesen K, *et al.* Segregation of lipid raft markers including CD133 in polarized human hematopoietic stem and progenitor cells. *Blood* 2004; 104: 2332-2338.
- [24] Bhatia M, Bonnet D, Murdoch B, Gan OI, Dick JE. A newly discovered class of human hematopoietic cells with SCID-repopulating activity. *Nat Med* 1998; 4: 1038-1045.
- [25] Zanjani ED, Almeida-Porada G, Livingston AG, Flake AW, Ogawa M. Human bone marrow CD34- cells engraft *in vivo* and undergo multilineage expression that includes giving rise to CD34+ cells. *Exp Hematol* 1998; 26: 353-360.
- [26] Gallacher L, Murdoch B, Wu DM, Karanu FN, Keeney M, Bhatia M. Isolation and characterization of human CD34-Lin- and CD34+Lin- hematopoietic stem cells using cell surface markers AC133 and CD7. *Blood* 2000; 95: 2813-2820.
- [27] Manz MG, Miyamoto T, Akashi K, Weissman IL. Prospective isolation of human clonogenic common myeloid progenitors. *Proc Natl Acad Sci U S A* 2002; 99: 11872-11877.
- [28] Galy A, Travis M, Cen D, Chen B, Human T, B, natural killer, and dendritic cells arise from a common bone marrow progenitor cell subset. *Immunity* 1995; 3: 459-473.
- [29] Moore MAS. The hematopoietic system and hematopoiesis. In *Neoplastic Hematopathology* (Knowels, D. M., Ed.), 2nd ed., pp. 1-42, Lippincott Williams & Wilkins, Philadelphia, PA; 2001.
- [30] Shah VO, Civin CI, Loken MR. Flow cytometric analysis of human bone marrow. IV. Differential quantitative expression of T-200 common leukocyte antigen during normal hemopoiesis. *J Immunol* 1988; 140: 1861-1867.
- [31] Terstappen LW, Safford M, Loken MR. Flow cytometric analysis of human bone marrow. III. Neutrophil maturation. *Leukemia* 1990; 4: 657-663.
- [32] Terstappen LW, Loken MR. Myeloid cell differentiation in normal marrow and acute leukemia assessed by multi-dimensional flow cytometry. *Anal Cell Pathol* 1990; 2: 229-240.
- [33] Clarke JL, Watkins W. Alpha1,3-L-fucosyltransferase expression in developing human myeloid cells. *J Biol Chem* 1996; 271: 10317-28.
- [34] Skacel PO, Edwards AJ, Harrison CT, Watkins WM. Enzymatic control of the expression of the X determinant (CD15) in human myeloid cells during maturation: The regulatory role of 6-sialyltransferase. *Blood* 1991; 78: 1452-1460.



- [35] Elghetany MT. Surface Antigen Changes during Normal Neutrophilic Development: A Critical Review. *Blood Cells Mol Dis* 2002; 28: 260-74.
- [36] Terstappen LW, Buescher S, Nguyen M, Reading C. Differentiation and maturation of growth factor expanded human hematopoietic progenitors assessed by multidimensional flow cytometry. *Leukemia* 1992; 6: 1001-1010.
- [37] De Jong MO, Wagemaker G, Wognum AW. Separation of myeloid and erythroid progenitors based on expression of CD34 and c-kit. *Blood* 1995; 86: 4076-4085.
- [38] Loken MR, Shah VO, Dattilio KL, Civin CI. Flow cytometric analysis of human bone marrow: I. Normal erythroid development. *Blood* 1987; 69: 255-263.
- [39] Rogers CE, Bradley MS, Palsson BO, Koller MR. Flow cytometric analysis of human bone marrow perfusion cultures: Erythroid development and relationship with burst-forming units-erythroid. *Exp Hematol* 1996; 24: 597-604.
- [40] Scicchitano MS, McFarland DC, Tierney LA, Narayanan PK, Schwartz LW. *In vitro* expansion of human cord blood CD36+ erythroid progenitors: Temporal changes in gene and protein expression. *Exp Hematol* 2003; 31: 760-769.
- [41] Debili N, Issaad C, Masse JM, Guichard J, Katz A, Breton-Gorius J, *et al.* Expression of CD34 and platelet glycoproteins during human megakaryocytic differentiation. *Blood* 1992; 80: 3022-3035.
- [42] Chang Y, Bluteau D, Debili N, Vainchenker W. From hematopoietic stem cells to platelets. *J Thromb Haemost* 2007; 1: 318-327.
- [43] Haddad R, Guardiola P, Izac B, Thibault C, Radich J, Delezoide AL, *et al.* Molecular characterization of early human T/NK and B-lymphoid progenitor cells in umbilical cord blood. *Blood* 2004; 104: 3918-3926.
- [44] Reynaud D, Lefort N, Manie E, Coulombel L, Levy Y. *In vitro* identification of human pro-B cells that give rise to macrophages, natural killer cells, and T cells. *Blood* 2003; 101: 4313-4321.
- [45] Hou YH, Srour EF, Ramsey H, Dahl R, Broxmeyer HE, Hromas R. Identification of a human B-cell/myeloid common progenitor by the absence of CXCR4. *Blood* 2005; 105: 3488-3492.
- [46] Akashi K, Traver D, Miyamoto T, Weissman IL. A clonogenic common myeloid progenitor that gives rise to all myeloid lineages. *Nature* 2000; 404: 193-197.
- [47] Katsura Y. Redefinition of lymphoid progenitors. *Nature Rev Immunol* 2002; 2: 127-132.
- [48] Lai AY, Kondo M. Asymmetrical lymphoid and myeloid lineage commitment in multipotent hematopoietic progenitors. *J Exp Med* 2006; 203: 1867-1873.
- [49] Ye M, Graf T. Early decisions in lymphoid development. *Curr Opin Immunol* 2007; 19: 123-128
- [50] Warren LA, Rothenberg EV. Regulatory coding of lymphoid lineage choice by hematopoietic transcription factors. *Curr Opin Immunol* 2003; 15: 166-175.
- [51] Ceredig R, Rolink AG, Brown G. Models of haematopoiesis: seeing the wood for the trees. *Nat Rev Immunol* 2009; 9: 293-300..

Received on 09-06-2014

Accepted on 26-06-2014

Published on 21-08-2014

© 2014 Armin Attar *et al.*; Licensee Cosmos Scholars Publishing House.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.