

Synergistic Antioxidant Activity of Capsicum Oleoresin, Lecithin and Curcuminoids in Sunflower Oil

Sameera A. Rege and Shamim A. Momin*

Department of Oils, Oleochemicals and Surfactants Technology, Institute of Chemical Technology, Nathalal Parekh Marg, Matunga (East), Mumbai – 400 019, India

Abstract: Essential fatty acids (EFAs) have the tendency to undergo autoxidation. Hence, the addition of antioxidants becomes a key step during their storage. Synthetic antioxidants are found to be toxic in nature. Recently, many investigations were carried out where EFAs were stabilized by natural antioxidants those can even replace synthetic antioxidant totally. In the present study, capsicum oleoresin, soy lecithin and curcuminoids were used to stabilize EFAs in refined sunflower oil. The study showed that the synergism of these natural antioxidants in proper proportion increased the shelf life of sunflower oil. Consequently, the nutritive value of EFAs was restored because of their effective stabilization and the natural antioxidants added for their stabilization provided the medicinal benefits.

Keywords: Antioxidants, Sunflower oil, Capsicum, Lecithin, Curcuminoids.

INTRODUCTION

Essential fatty acids (EFAs) are required dietary nutrients as they perform vital functions for good health and cannot be synthesized completely in human body [1]. The main sources of these EFAs are oils such as flaxseed, soybean, sunflower, safflower and fish oils [2]. Being highly susceptible to autoxidation, the storage of these oils becomes difficult. The process of autoxidation needs to be prevented to restore the nutritive value of EFAs. Generally, synthetic antioxidants such as tertiary butylhydroquinone (TBHQ), butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) are added to prevent the autoxidation but their safety has been questioned [3]. Accordingly, a safer antioxidant needs to be investigated.

Recent research has been concentrated on natural sources of antioxidants such as spices, herbs [4]. The presence of active components in spices has been demonstrated over the last 30 years [5]. Spices and condiments, though employed in small quantities, exert a considerable effect on the stability of fats and oils [6, 7]. Hence, it is worth to state that the use of these natural antioxidants is a promising substitute to the synthetic antioxidants.

The common household spices extensively used in India, capsicum [8] and curcuminoids [9] are known to have antioxidant effect. The capsicum was found to be

effective antioxidant in oil [9] and margarine as well [10]. Herbalists have valued capsicum for its various therapeutic actions including anti-inflammatory, antiseptic effects [11], treatment of cutaneous allergy and neurological disorders [12]. Emerging studies have shown that capsicum is an effective medicine against neuropathic pain [13]. Capsaicinoids may have the potential for the weight loss [14] and also as an anticancer agent [15]. Similarly, curcumin, a major component of curcuminoids has medicinal properties against various diseases such as anorexia, coughs, diabetes, hepatic disorders, rheumatism and Alzheimer disease [16]. Curcuminoids are phytopharmaceuticals that have been shown to alleviate depressive symptoms [17] and an effective agent against wound healing [18].

It has been reported previously that curcuminoids alone revealed pro-oxidant effect [19]. However, the addition of these antioxidants in a suitable concentration can exert an antioxidant effect as a whole. The present study focuses to evaluate the antioxidant activity of capsicum oleoresin alone and in synergism with curcuminoids in sunflower oil. The work was further extended to study the synergistic activity of capsicum oleoresin, curcuminoids and soy lecithin, a well known antioxidant [20, 21] in the same oil.

MATERIALS AND METHODS

Materials

Refined sunflower oil without added antioxidant (RSFO) was received as a gift sample from M/s Cargill India Pvt. Ltd., New Delhi. The oil did not contain any added synthetic antioxidant. Soy lecithin was procured

*Address correspondence to this author at the Department of Oils, Oleochemicals and Surfactants Technology, Institute of Chemical Technology, Nathalal Parekh Marg, Matunga (East), Mumbai – 400 019, India; Tel: +91-9870148606; E-mail: samomin@rediffmail.com

from M/s V.R. Chemicals, Mumbai. Capsicum oleoresin and curcuminoids powder were obtained from Kancor India Ltd., Angamaly South, India. All other chemical reagents and solvents were obtained from s.d. fiNE-CHEM LiMiTEd, Mumbai.

Analysis of Capsicum Oleoresin

The presence of capsaicinoids was checked by UV spectrometry (Shimadzu UV-1601) of capsicum oleoresin in isopropanol [22].

Preparation of Samples to Study Antioxidant Activity

The various blends of oil containing capsicum oleoresin (1.5%, 2%, 2.5%, 3% w/v) and its combination with soy lecithin (1.5% w/v) and curcuminoids (50 ppm) were prepared. The oxidative stability of oil blends was checked at 60°C for 30 days at regular interval of 5 days according to the AOCS Official Methods [23] by peroxide value (PV, Method Cd 8-53), *p*-anisidine value (*p*-A.V., Method Cd 18-90) and total oxidation (Totox) value (Method Cg 3-91). All the experiments were carried out in triplicate and the values were expressed as arithmetic mean of the experiments along with standard deviation.

The activities of natural antioxidants were compared with control sample and TBHQ (200 ppm) under the same conditions. The relative antioxidant activities were compared using Oxidative Factor (OXF) for antioxidants based on mean peroxide value of triplicate experiments using following formula [24],

$$OXF = \frac{(PV_{final} - PV_{initial})_{antioxidant}}{(PV_{final} - PV_{initial})_{control}}$$

where PVs indicate the mean values of all triplicate determinations of the peroxide value.

RESULTS AND DISCUSSION

Linoleic and linolenic acids are respectively 10 and 25 times more unstable as compared to monounsaturated oleic acid [25]. The deterioration of sunflower oil rich in linoleic acid can be prevented by using a suitable antioxidant. The antioxidants are phenolic compounds substituted by electron donating groups at *ortho* and/or *para* positions. They act by inactivating free radicals generated during autoxidation process by proton donation mechanisms [26]. In this

context, easily available as well as non-toxic natural antioxidants like capsicum oleoresin, soy lecithin and curcuminoids were used as substitutes for synthetic antioxidant.

The presence of capsaicinoids in capsicum oleoresin was confirmed by the peak at 281 nm in the UV spectrum [22] of the capsicum oleoresin solution in isopropanol.

Oxidative Stability of Sunflower Oil in Presence of Capsicum Oleoresin

The peroxides are primary oxidation products that can further undergo degradation to form low molecular weight aldehydes and ketones. The radicals are prone to peroxidation and follow the chain reactions, typical of radicals [25]. Capsicum oleoresin contains a group of capsaicinoids like capsaicin, dihydrocapsaicin, nordihydrocapsaicin, homocapsaicin, homodihydrocapsaicin that act as conventional antioxidants by deactivating radicals. Semwal *et al.* [27] showed that capsaicin and dihydrocapsaicin reveal considerable antioxygenic activity when incorporated at the concentration of 100 ppm in sunflower oil. The solubility of capsicum oleoresin in RSFO was found to be 3%. The capsicum oleoresin under study contained 3.14% capsaicinoids. Consequently, capsicum oleoresin did not show significant antioxidant activity at lower concentration (200-800 ppm) since it did not provide sufficient amount of capsaicinoids. It showed significant antioxidant activity (Tables 1, 2 and Figure 1) at higher concentrations (2-3%) corresponding to 600-1000 ppm capsaicinoids. There was insignificant increase in antioxidant activity with increase in the concentration beyond 2.5% concentration.

Effect of Capsicum Oleoresin (2.5%), Lecithin and Curcuminoids on Oxidative Stability of RSFO

Curcuminoids exhibit keto-enol tautomerism in solutions, with up to 95% curcuminoids in the enol form depending on the solvent [28]. In alkaline conditions, curcuminoids rapidly decompose to ferulic acid and feruloylmethane [29] that further undergo hydrolysis to form vanillin and acetone. Masuda *et al.* [30] also supported autoxidation of curcuminoids. In neutral and acidic solutions (from pH 3 to 7), the keto form predominates that acts as H-atom donor. Thus, it is crucial to maintain these curcuminoids in the keto form by employing acidic conditions to utilize their antioxidant activity.

Initially, capsicum oleoresin (2.5%) was incapable to stabilize curcuminoids thereby showing peroxide formation same as that of RSFO (Tables 1, 2 and Figure 2). But as the study was continued, it showed marginal antioxidant activity in synergism with curcuminoids. However, it was lower than that of capsicum oleoresin alone. Though capsicum is a good source of phenolic acids [31], the amount was insufficient to stabilize curcuminoids. On the other hand, the same blend showed lower totox value. It indicated that the blend has potential to stabilize peroxy radical thereby reducing the formation of secondary degradation products like carbonyl compounds.

Capsicum oleoresin showed better antioxidant activity in synergism with lecithin. Furthermore, when the mixture of curcuminoids is added to the same blend, the antioxidant activity was improved. It was comparable to the activity of TBHQ (Figure 2). Lecithin is known to prevent the autoxidation of oil by its action as an oxygen barrier between oil/air interface and also as a metal ion chelator [32]. The radical deactivating potential of capsaicinoids was accompanied by oxygen barrier from lecithin resulting in reduced peroxide formation.

The presence of free fatty acids from lecithin (acid value, 6.23 mg.g⁻¹) inhibited the degradation of

Table 1: Effect of Synergism of Capsicum Oleoresin with Lecithin and Curcuminoids on Oxidative Stability (Peroxide Formation) of Refined Sunflower Oil without Antioxidant (RSFO) at 60°C

Sample	Time (days)	OXF					
		5	10	15	20	25	30
RSFO + TBHQ		0.66	0.40	0.28	0.28	0.21	0.32
RSFO + 2% capsicum oleoresin		0.44	0.87	0.61	0.72	0.60	0.71
RSFO + 2.5% capsicum oleoresin		0.51	0.50	0.52	0.58	0.56	0.66
RSFO + 3% capsicum oleoresin		0.56	0.48	0.51	0.58	0.52	0.59
RSFO + 2.5% capsicum oleoresin + 50 ppm curcuminoids		1.16	1.04	0.85	0.75	0.79	0.72
RSFO + 2.5% capsicum oleoresin + 1.5% lecithin		0.83	0.63	0.63	0.51	0.47	0.47
RSFO + 1.5% capsicum oleoresin + 1.5% lecithin + 50 ppm curcuminoids		1.21	0.85	0.82	0.78	0.80	0.75
RSFO + 2% capsicum oleoresin + 1.5% lecithin + 50 ppm curcuminoids		1.02	0.76	0.74	0.64	0.52	0.62
RSFO + 2.5% capsicum oleoresin + 1.5% lecithin + 50 ppm curcuminoids		1.17	0.70	0.48	0.44	0.39	0.30

Table 2: Effect of Synergism of Capsicum Oleoresin with Lecithin and Curcuminoids on Oxidative Stability (Totox Value in meq/kg) of Refined Sunflower Oil without Antioxidant (RSFO) at 60°C

Sample	Time (days)	TOTOX [†]					
		5	10	15	20	25	30
RSFO		17.0±1.4	22.6±1.5	30.0±1.5	40.1±1.6	51.5±2.2	70.9±3.9
RSFO + TBHQ		6.3±0.9	8.2±0.8	11.2±1.0	14.0±1.1	14.5±1.6	26.3±2.8
RSFO + 2% capsicum oleoresin		11.1±1.0	15.0±1.0	19.8±0.9	29.1±1.6	33.3±1.7	51.0±2.5
RSFO + 2.5% capsicum oleoresin		9.2±1.2	12.7±1.2	17.5±1.3	24.8±1.5	30.5±1.7	47.1±2.4
RSFO + 3% capsicum oleoresin		9.3±1.1	12.5±1.3	16.9±0.7	23.8±1.2	28.9±1.4	43.2±2.2
RSFO + 2.5% capsicum oleoresin + 50 ppm curcuminoids		9.6±1.5	12.7±1.2	18.0±1.0	20.2±1.2	29.4±1.5	40.4±2.2
RSFO + 2.5% capsicum oleoresin + 1.5% lecithin		10.5±0.8	13.6±1.0	19.0±1.2	21.0±2.0	29.4±2.2	40.9±3.4
RSFO + 1.5% capsicum oleoresin + 1.5% lecithin + 50 ppm curcuminoids		12.8±0.7	15.2±1.2	17.6±0.9	21.7±1.1	29.6±1.6	40.9±1.8
RSFO + 2% capsicum oleoresin + 1.5% lecithin + 50 ppm curcuminoids		8.7±1.0	11.8±0.7	16.6±0.6	20.6±1.1	26.7±1.5	33.2±1.9
RSFO + 2.5% capsicum oleoresin + 1.5% lecithin + 50 ppm curcuminoids		8.1±0.6	10.4±0.9	14.7±0.5	16.6±1.3	17.8±1.6	29.1±2.2

[†] The values given are means of three consecutive experiments ± standard deviations




Figure 1: Effect of capsicum oleoresin on oxidative stability (peroxide formation) of RSFO at 60°C.




Figure 2: Effect of capsicum oleoresin, lecithin and curcuminoids on oxidative stability (peroxide formation) of RSFO at 60°C.

curcuminoids thereby maintaining them in keto form. A similar mechanism was shown by kalonji seeds extract as well that also inhibited the degradation of curcuminoids due to presence of free fatty acids [33]. As a result, lecithin, capsicum oleoresin and

curcuminoids all acted by different mode of action and concurrently showed synergistic antioxidant activity. In addition, there was increase in antioxidant activity with increase in the concentration of capsicum oleoresin in presence of lecithin and curcuminoids (Tables 1 and 2).

CONCLUSION

The synergistic antioxidant activity of capsicum oleoresin, lecithin and curcuminoids in appropriate quantity in refined sunflower oil was comparable to synthetic antioxidant. This effect was due to the combined action of the added antioxidants resulting in increased antioxidant potential than that expected from a mere additive effect. Natural antioxidants being the choice of consumers, can find better acceptability over synthetic antioxidants. With stabilization of essential fatty acids (EFAs) using such antioxidants in conjunction with their medicinal benefits, the consumers can enjoy the benefits of EFAs and therapeutic uses of natural antioxidants. Since these natural extracts containing antioxidants are normally used in household cooking, their taste will be familiar and hence acceptable.

ACKNOWLEDGEMENTS

This study was financially supported by Technical Education Quality Improvement Programme (TEQIP), Government of India and World Bank.

REFERENCES

- [1] Spector AA. Essentiality of fatty acids. *Lipids* 1999; 34: S1-S3. <https://doi.org/10.1007/BF02562220>
- [2] Orthoefer FT. Vegetable oils. In: Hui YH, editor. *Bailey's Industrial Oils & Fats, Vol.1. Edible oil & fat products: General applications*. 5th ed. New York: John Wiley & Sons Inc. 1996; p. 19-43.
- [3] Barlow SN. In: Hudson BJB, editor. *Toxicological Aspects of Antioxidants Used as Food Additives*. Food Antioxidants. Amsterdam: Elsevier 1990; p. 253-307.
- [4] Suhaj M. Spice antioxidants isolation and their antiradical activity: A review. *J Food Composition Anal* 2006; 19: 531-537. <https://doi.org/10.1016/j.jfca.2004.11.005>
- [5] Lagouri V, Boskou D. Screening for antioxidant activity of essential oils obtained from spices. *Dev Food Sci* 1995; 37: 869-879. [https://doi.org/10.1016/S0167-4501\(06\)80203-7](https://doi.org/10.1016/S0167-4501(06)80203-7)
- [6] Kim SY, Kim JH, Kim SK, Oh MJ, Jung MY. Antioxidant activities of selected oriental herb extracts. *J Am Oil Chem Soc* 1994; 71: 633-640. <https://doi.org/10.1007/BF02540592>
- [7] Patki PE, Arya SS. Studies on development and storage stability of instant dhals. *Indian Food Packer* 1994; 48: 31-39.
- [8] Nag A. Stabilization of flaxseed oil with capsicum antioxidant. *J Am Oil Chem Soc* 2000; 77: 799-800. <https://doi.org/10.1007/s11746-000-0127-0>
- [9] Kunchandy E, Rao MNA. Oxygen radical scavenging activity of curcumin. *Int J Pharm* 1990; 58: 237-240. [https://doi.org/10.1016/0378-5173\(90\)90201-E](https://doi.org/10.1016/0378-5173(90)90201-E)
- [10] Rege SA, Momin SA, Wadekar SD, Bhowmick DN. Formulation of a functional fat spread stabilized by natural antioxidants and emulsifiers *Malays J Nutr* 2013; 19: 121-129.
- [11] Dasgupta P, Fowler CJ. Chillies: from antiquity to urology. *Br J Urol* 1997; 80: 845-852. <https://doi.org/10.1046/j.1464-410X.1997.00424.x>
- [12] Palevitch D, Craker LE. Nutritional and medicinal importance of red pepper (*Capsicum* spp.). *J Herbs Spices Med Plants* 1995; 3: 55-83. https://doi.org/10.1300/J044v03n02_08
- [13] Yong YL, Tan LT, Ming LC, Chan KG, Lee LH, Goh BH, et al. The effectiveness and safety of topical capsaicin in postherpetic neuralgia: A systematic review and meta-analysis. *Front Pharmacol* 2017; 7: 538. <https://doi.org/10.3389/fphar.2016.00538>
- [14] Srinivasan K. Biological activities of red pepper (*Capsicum annum*) and its pungent Principle Capsaicin: A Review. *Crit Rev Food Sci Nutr* 2016; 56: 1488-1500. <https://doi.org/10.1080/10408398.2013.772090>
- [15] Shaimaa GA, Mahmoud MS, Mohamed MR, Emam AA. Phytochemical screening, antioxidant activities and in vitro anticancer potential of Egyptian *Capsicum* Spp. *Biochem Pharmacol* 2016; 5:2.
- [16] Wang X, Jiang Y, Wang YW, Huang MT, Ho CT, Huang Q. Enhancing anti-inflammation activity of curcumin through O/W nanoemulsions. *Food Chem* 2008; 108: 419-424. <https://doi.org/10.1016/j.foodchem.2007.10.086>
- [17] Panahi Y, Badeli R, Karami GR, Sahebkar A. Investigation of the efficacy of adjunctive therapy with bioavailability-boosted curcuminoids in major depressive disorder. *Phytother Res* 2015; 29: 17-21. <https://doi.org/10.1002/ptr.5211>
- [18] Kant V, Gopal A, Pathak NN, Kumar P, Tandan SK, Kumar D. Antioxidant and anti-inflammatory potential of curcumin accelerated the cutaneous wound healing in streptozotocin-induced diabetic rats. *Int Immunopharmacol* 2014; 20: 322-330. <https://doi.org/10.1016/j.intimp.2014.03.009>
- [19] Semwal AD, Sharma GK, Arya SS. Antioxygenic activity of turmeric (*Curcuma longa*) in sunflower oil and ghee. *J Food Sci Tech* 1997; 34: 67-69.
- [20] Hildebrand DH, Terao J, Kito M. Phospholipids plus tocopherols increase soybean oil stability. *J Am Oil Chem Soc* 1984; 61: 552-555. <https://doi.org/10.1007/BF02677029>
- [21] Jung U, Lee SK, Kyung SH, Chang GH, Chang IM, Lee CH. Antioxidant effects of natural lecithin on borage oil. *Food Sci Biotechnol* 2001; 10: 354-359.
- [22] Gonzalez AT, Tamirano CW. A new method for the determination of capsaicin in *Capsicum* fruits. *J Food Sci* 1973; 38: 342-345. <https://doi.org/10.1111/j.1365-2621.1973.tb01422.x>
- [23] AOCS: 1994. Firestone D, editor. *Official methods and recommended practices of the American Oil Chemists' Society*. 4th ed. Champaign, IL: Am. Oil Chem. Soc. 1994.
- [24] Ghada A, Vassiliki O. Antioxidant properties and composition of Majorana syriaca extracts. *Eur J Lipid Sci Technol* 2007; 109: 247-255. <https://doi.org/10.1002/ejlt.200600234>
- [25] Knothe G, Kenar JA, Gunstone FD. Chemical Properties. In: Gunstone FD, Harwood JL, Dijkstra AJ, editors. *The Lipid Handbook*. LLC: Taylor & Francis Group 2007; p. 535-541.
- [26] Khan MA, Shahidi F. Effects of natural and synthetic antioxidants on the oxidative stability of Borage and Evening Primrose Triacylglycerols. *Food Chem* 2001; 75: 431-437. [https://doi.org/10.1016/S0308-8146\(01\)00232-1](https://doi.org/10.1016/S0308-8146(01)00232-1)
- [27] Semwal AD, Sharma GK, Arya SS. Pro- or antioxygenic activity of tejpatt (*Cinnamomum tamala*) and red chili (*Capsicum annum*) in sunflower oil. *J Sci Food Agric* 1999; 79: 1733-1736. [https://doi.org/10.1002/\(SICI\)1097-0010\(199909\)79:12<1733::AID-JSFA431>3.0.CO;2-4](https://doi.org/10.1002/(SICI)1097-0010(199909)79:12<1733::AID-JSFA431>3.0.CO;2-4)

- [28] Chemical and Technical Assessment, 61st JECFA, 2004. Curcumin [cited 2017 Jan 26]: Available from: <ftp://ftp.fao.org/esn/jecfa/cta/CTA_61_Curcumin.pdf>.
- [29] Tonnesen HH, Karlsen J. Studies of curcumin and curcuminoids: V. Alkaline degradation of curcumin. *Z Lebensm Unters Forsch* 1985; 180: 132-134. <https://doi.org/10.1007/BF01042637>
- [30] Masuda T, Hidaka K, Shinohara A, Maekawa T, Takeda Y. Chemical studies on antioxidant mechanism of curcuminoid: Analysis of radical reaction products from curcumin. *J Agric Food Chem* 1999; 47: 71-77. <https://doi.org/10.1021/jf9805348>
- [31] Benzie IF. Evolution of dietary antioxidants. *Comp Biochem Physiol Part A: Mol Integr Physiol* 2003; 136: 113-126. [https://doi.org/10.1016/S1095-6433\(02\)00368-9](https://doi.org/10.1016/S1095-6433(02)00368-9)
- [32] Hudson B, Ghavani M. Phospholipids as antioxidant synergists for tocopherols in the autoxidation of edible oils. *Lebensm-Wiss. Technol* 1984; 17: 191-194.
- [33] Rege SA, Momin SA, Bhowmick DN, Pratap AP. Stabilization of emulsion and butter like products containing essential fatty acids using kalonji seeds extract and curcuminoids. *J Oleo Sci* 2012; 61: 11-16. <https://doi.org/10.5650/jos.61.11>

Received on 26-01-2017

Accepted on 07-02-2017

Published on 17-02-2017

<http://dx.doi.org/10.15379/2408-9826.2017.04.01.02>

© 2017 Rege and Momin; Licensee Cosmos Scholars Publishing House.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.